

# **Autonomous Benthic Monitoring Downunder**

O. Pizarro, S.B. Williams, M. Bryson, A. Friedman





### **Integrated Marine Observing System**








#### **Sirius AUV**

- Flexible, mobile, high resolution data collection
- Sensors include
  - DVL
  - Attitude
  - Pressure depth
  - fwd obstacle avoidance
  - GPS (surface)
  - USBL
  - Vision (stereo)
  - Multibeam sonar
  - Water Column












# **SLAM: Visually Augmented Navigation**











### **IMOS** benthic monitoring program









#### **Australia's Network of Marine Protected Areas**


















### **Cyclone Ita path**














### Involute of a circle

















# Lizard Island post cyclone surveys









### Path Length, area and spacing design curve












Slide 1199


















#### 'Reef Records'

- High overlap by design
- Robust to swell and current
- Easy line handling underwater
- Monitoring through revisiting sites
- Visible range, 6m radius
- •~110m<sup>2</sup> area

	Avg.	Std. Dev.	Max.	Min.
Duration	15:26	01:43	18:47	12:58
Stereo pairs	1853	205.3	2255	1557
Image matches	19286.3	7098.1	33168	6194



# Lizard Island - post cyclones ITA (and Nathan)

Lizard Island


National Park

- APR 2014: 7 days, 23 sites
- OCT 2014:7 days, 23 sites
- APR 2015:5 days, 21+ sites
- NOV 2015, 21+ sites
- Data processed and delivered in field
- NOV 2016, 21+ sites






# **Structural Complexity Measurement:** Rugosity



Friedman et. al. 2013, Multi-scale measures of rugosity, slope and aspect from benthic stereo image reconstructions




# **Structural Complexity Measurement:** Rugosity




Friedman et. al. 2013, Multi-scale measures of rugosity, slope and aspect from benthic stereo image reconstructions




#### **Real Chain to Virtual chain**














Horseshoe reefrecord Repeated OCT 2014













# ACFR Horseshoe reefrecord repeats








# Results: distribution of rugosity errors for each specific survey time






- For both single day and multiple day surveys, some sites exhibited significant per-survey biases (up to 7.5% of rugosity range)
- Highlights the importance in changes in survey conditions and survey trajectory parameters having a large influence on the actual measurements being made






# SC measurement errors vs. mean rugosity








# SC measurement errors vs. mean rugosity and coral morphotype
























April 2010



# April 2011






































### **Acknowledgements**



- The technical staff of the ACFR make all this possible. As special thanks to Christian Lees, Andy Durrant and Jeremy Randle.
- Our science collaborators, particularly Josh Madin, Renata Ferrari, Gary Kendrick and Maria Dornelas
- This IMOS AUV facility is supported in part by NCRIS IMOS, the Australian Research Council (ARC) and the New South Wales Government
- Additional funding has been received from our industry and government partners, including SEA SERPENT







